arXiv:1212.6809v1 [math.OA] 31 Dec 2012

About the bound of the C* exponential length

Qingfei Pan and Kun Wang

December, 2012

Abstract

In this paper, we give an example to show that, if $u \in \mathrm{C}(X)\otimes\mathrm{M}_n$ with $\det(u) = 1$ then the C* exponential length of u (denoted by $cel(u)$) can not be controlled by π. Moreover, in the simple inductive limit C*-algebras, similar examples exist.

Introduction. The C* exponential length, as a related invariant of C* exponential rank, has been studied for many years. In [8], N.C. Phillips mentioned that "Suitable modifications of Lemma 5.2 and 5.3 will show that if $\det(u) = 1$, then $cel(u) \leq \pi$ (even though, for general u, $cel(u)$ can be arbitrarily large)." But this is not true. For any $\varepsilon > 0$, we can find a unitary u in a C*-algebra with $\det(u) = 1$ and $cel(u) \geq 2\pi - \varepsilon$. In this paper, we provide a method for constructing such examples. And in the future we will show that, under the same assumption, 2π is an upper bound.

1. Preliminaries.

For convenience of the reader, we list following definitions and lemmas (see [3], [10] for more details).

Definition 1.1. Let X be a compact space and $B = \mathrm{C}(X)\otimes\mathrm{M}_n$, for $u \in U(B)$(unitary group of B), let $\det(u)$ be a function from x to S^1 whose value at x is $\det(u(x))$.

The following lemma is well-known.

Proposition 1.2. Let u be a unitary in a C*-algebra A and suppose $u \sim_h 1$, then

$$u = \exp(ih_1)\exp(ih_2)...\exp(ih_k),$$

for some self-adjoint elements h_i's in A and some integer k.

Definition 1.3. Let A be a unital C*-algebra and u be a unitary which lies in the connected component of the identity 1 in A. Define the C* exponential length of u (denoted by $cel(u)$) as follows:

$$cel(u) = \inf\{\sum_{i=1}^{k} \|h_i\| : u = \exp(ih_1)\exp(ih_2)...\exp(ih_k)\}.$$

Note: For a unital C*-algebra A let $U_0(A)$ be the connected component of $U(A)$ containing the identity 1. Recall from [10] that if $u \in U_0(A)$, then the C* exponential length $cel(u)$ is equal to the infimum of the lengths of paths in $U(A)$ from u to 1.

Following lemma is an easy example for calculating the C* exponential length.

Lemma 1.4. Let $u(t) = \exp(itA)$ be a unitary in C$[0,1]$ with A a real number, then

$$cel(u) = |A|.$$

Proof: Since $cel(u) = \inf\{\text{length}(u_s) : u_s \text{ is a path in } U(A) \text{ from } u \text{ to } 1\}$, let $v_s(t)$ be any path from 1 to u, that is, $v_0(t) = 1, v_1(t) = u(t)$. Then length$(v_s) = \int_0^1 \|\frac{dv}{ds}\|ds$. Since $v_s(t)$ can be considered as a map from $[0,1] \times [0,1]$ to S^1 and $\mathbb{R}$ is a covering space of S^1, there exists $\tilde{v}$ from $[0,1] \times [0,1]$ to $\mathbb{R}$ such that:

$$v(s,t) = \pi(\tilde{v}(s,t)) \text{ and } \widetilde{v}((0,t)) = 0,$$

where $\pi(x) = e^{ix}$. Therefore

$$\frac{dv}{ds} = \pi'(\tilde{v}(s,t)) \cdot \frac{d\widetilde{v}}{ds}, \quad \|\frac{dv}{ds}\| = \|\frac{d\tilde{v}}{ds}\|.$$

Since $\tilde{v}$ maps $(0,t)$ to 0, $(1,t)$ to tA,

$$\int_0^1 \|\frac{d\tilde{v}}{ds}\|ds \geq |\int_0^1 \frac{d\tilde{v}}{ds}ds| = \max_t |v(1,t) - v(0,t)| = |A|.$$

On the other hand, $\exp(istA)$ for $0 \leq s \leq 1$ is a path in $U(\text{C}[0,1])$ from 1 to u with length $|A|$. Therefore $cel(u) = |A|$.

2. Counter Examples.

Lemma 2.1.([4] Lemma 2.5). Let X be a finite simplicial complex of dimension at most 2. Let E be a locally trivial M_n-bundle over X, let $u \in \Gamma(SU_E)$ and let $\varepsilon > 0$. Then there exists $v \in \Gamma(SU_E)$ such that $\|u - v\| < \varepsilon$ and $v(x)$ has no repeated eigenvalues for all $x \in X$, where SU_E is the set of elements of E which are unitaries with determinant 1 and $\Gamma(U_E)$ is the set of sections of U_E.

Corollary 2.2. Let u_s $(0 \leq s \leq 1)$ be a path in $U(\text{M}_{10}(\text{C}[0,1]))$ and let $\varepsilon > 0$, then there exists v_s $(0 \leq s \leq 1)$ a path in $U(\text{M}_{10}(\text{C}[0,1]))$ such that $\|u - v\| < \varepsilon$ and $v_s(t)$ has no repeated eigenvalues for all $0 \leq s \leq 1$ and $0 \leq t \leq 1$.

Proof: By considering $u_s(t)$ as a map from $[0,1] \times [0,1]$ to $U(\text{M}_{10}(\mathbb{C}))$, the Corollary follows from the Theorem 2.1.

Let $P^kS^1 := \{[x_1, x_2, ..., x_k] : x_i \in S^1\}/\sim$, where $[x_1, x_2, ..., x_k] \sim [y_1, y_2, ..., y_k]$ if $\exists \sigma \in S_k$ such that $y_{\sigma(i)} = x_i$. Naturally, the metric of P^kS^1 should be defined as:

$$\text{dist}([x_1, x_2, ..., x_k], [y_1, y_2, ..., y_k]) = \min_{\sigma} \max_{1 \leq i \leq k} \text{d}(x_i, y_{\sigma(i)}).$$

The following lemma is well-known:

Lemma 2.3. Let $F : [0,1] \times [0,1] \to P^kS^1$ be a continuous function, suppose

$$F(s,t) = [x_1(s,t), x_2(s,t), ..., x_k(s,t)],$$

and for any fixed $(s,t) \in [0,1] \times [0,1]$, $x_i(s,t)$ $(1 \le i \le k)$ are k distinct values in S^1. Then there exist $f_1, f_2, ..., f_k : [0,1] \times [0,1] \to S^1$ such that:

$$F(s,t) = [f_1(s,t), f_2(s,t), ..., f_k(s,t)].$$

Proof: Since F is continuous, $F([0,1] \times [0,1])$ is a connected set in P^kS^1. Let $\pi : \underbrace{S^1 \times S^1 \times ... \times S^1}_{k} \to P^kS^1$ denote the quotient map, and let $X \subset \underbrace{S^1 \times S^1 \times ... \times S^1}_{k}$ consists of those elements $(x_1, x_2, \cdots, x_k)$ with $x_i \neq x_j$ for $i \neq j$. Then $\pi|_X$ is a covering map from X to $\pi(X)$ (which is a subset of P^kS^1).

Note from the condition of the Lemma, the image of the map $F : [0,1] \times [0,1] \to P^kS^1$ is inside $\pi(X)$. Since $[0,1] \times [0,1]$ is simply connected, by standard lifting theorem for covering spaces, the map $F : [0,1] \times [0,1] \to \pi(X) \subset P^kS^1$ can be lifted to a map $F_1 : [0,1] \times [0,1] \to X(\subset \underbrace{S^1 \times S^1 \times ... \times S^1}_{k})$

Let $\pi_j : S^1 \times S^1 \times ... \times S^1 \to S^1$ be the projection onto the jth coordinate. For $1 \le j \le k$, define functions $f_j : [0,1] \times [0,1] \to S^1$ by

$$f_j(s,t) = \pi_j F_1(s,t).$$

Then it is easy to see that f_j's satisfy the requirement.

Example 2.4. Let $A = \mathrm{M}_{10}(\mathrm{C}[0,1])$, define

$$u(t) = \begin{pmatrix} e^{-2\pi i t \frac{9}{10}} & 0 & \cdots & 0 \\ 0 & e^{it\frac{2\pi}{10}} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & e^{it\frac{2\pi}{10}} \end{pmatrix}_{10 \times 10}.$$

Then u is a unitary in A with $\det(u) = 1$ and $u \sim_h 1$.

Theorem 2.5. Let $u \in \mathrm{M}_{10}(\mathrm{C}[0,1])$ be defined as in 2.4, then

$$cel(u) \ge \frac{9}{10} \cdot 2\pi.$$

Proof: Let $\widetilde{F}_s$ be a path in $U(\mathrm{M}_{10}(\mathrm{C}[0,1]))$ with $\widetilde{F}_0 = 1 \in \mathrm{M}_{10}(\mathrm{C}[0,1])$ and $\widetilde{F}_1(t) = u(t)$.

By Corollary 2.2 above, there is a path F_s in $U(\mathrm{M}_{10}(\mathrm{C}[0,1]))$ such that $\|F-\widetilde{F}\| < \varepsilon/2$ and $F_s(t)$ has no repeated eigenvalues for all $s \in [0,1]$ and $t \in [0,1]$. By Lemma 2.3, there exist $f^1, f^2, ..., f^k : [0,1] \times [0,1] \to S^1$ and unitaries $U_s(t)$ such that $F_s(t) = U_s(t)[f^1_s(t), f^2_s(t), ..., f^k_s(t)]U_s(t)^*$. Therefore

$$\mathrm{length}(F_s) = \max_{1\le i\le k}\{\mathrm{length}(f^i_s)\}.$$

Since $\|F-\widetilde{F}\| < \varepsilon/2$, $\|f^j_0 - 1\| < \varepsilon/2$ for all $1 \le j \le k$. For each fixed $t \in (\varepsilon, 1-\varepsilon)$, there exists one and only one j_0 such that

$$\|f^{j_0}_1(t) - e^{-2\pi i t\frac{9}{10}}\| < \varepsilon/2$$

and for other $j \neq j_0$

$$\|f^j_1(t) - e^{it\frac{2\pi}{10}}\| < \varepsilon/2.$$

(We use the fact that $\|e^{it\frac{2\pi}{10}} - e^{-2\pi i t\frac{9}{10}}\| \ge \varepsilon$ for $t \in (\varepsilon, 1-\varepsilon)$.) Since all f^j's are continuous, the index j_0 should be the same for all $t \in (\varepsilon, 1-\varepsilon)$.

Therefore, $f^{j_0}_s$ is a path in $U(\mathrm{C}[0,1])$ connecting a point near 1 and a point near $e^{-2\pi i t\frac{9}{10}}$. By Lemma 1.4, $\mathrm{length}(f^{j_0}_s) \ge \frac{9}{10}\cdot 2\pi - \varepsilon$. Then,

$$\mathrm{length}(\tilde{F}_s) \ge \frac{9}{10}\cdot 2\pi - \varepsilon/2 - \varepsilon \ge \frac{9}{10}\cdot 2\pi - 2\varepsilon.$$

Since ε is arbitrary, we have $cel(u) \ge \frac{9}{10}\cdot 2\pi$, which completes the proof.

Example 2.6. In some simple inductive limit C*-algebras, such examples exist.
Let $\{x_1, x_2, ...\}$ be a countable dense subset of $[0,1]$, and let $\{k_n\}_{n=2}^{\infty}$ be a sequence of integers satisfying

$$\prod_n \frac{10^{k_n}-1}{10^{k_n}} > \frac{7}{8}.$$

Let

$$A_1 = \mathrm{M}_{10}(\mathrm{C}[0,1]),\ A_2 = \mathrm{M}_{10^{k_2}}(A_1),\ ...,\ A_n = \mathrm{M}_{10^{k_n}}(A_{n-1}), ...$$

Define $\varphi_{n,n+1} : A_n \to A_{n+1}$ by

$$\varphi_{n,n+1}(f) = \begin{pmatrix} f & & & \\ & \ddots_{\;10^{k_n}-1} & & \\ & & f & \\ & & & f(x_n) \end{pmatrix}.$$

Let $A = \varinjlim(A_i, \varphi_{i,i+1})$ be the inductive limit C*-algebra. Then A is simple.
Let $u(t) \in A_1$ be defined as in Example 2.4, then a similar way of Theorem 2.5 shows that

$$cel(\varphi_{1,\infty}(u)) \ge \frac{9}{10}\cdot 2\pi.$$

References

[1] M.-D. Choi and G.A. Elliott, *Density of the self-adjoint elements with finite spectrum in an irrational rotation C*-algebra*, Math. Scand. 67(1990), 73-86.

[2] V. Guillemin and A. Pollack, *Differential Topology.*

[3] R.V. Kadison and J.R. Ringrose, *Fundamentals of the theory of operator algebras*, vol. 1, Elementary theory (New York: Academic Press, 1983).

[4] N.C. Phillips, *Simple C*-algebras with the property weak (FU)*, Math. Scand. 69(1991), 127-151.

[5] N.C. Phillips, *The rectifiable metric on the space of projections in a C*-algebra*, Internat. J. Math. 3(1992), 679-698.

[6] N.C. Phillips, *How many exponentials?*, Amer. J. Math., Vol. 116, No. 6 (Dec., 1994), 1513-1543.

[7] N.C. Phillips, *The C* projective length of n-homogeneous C*-algebras*, J. Operator Theory, 31(1994), 253-276.

[8] N.C. Phillips, *Reduction of exponential rank in direct limits of C*-algebras*, Can. J. Math. Vol. 46(4), 1994, 818-853.

[9] N.C. Phillips, *Exponential length and traces*, Proc. Roy. Soc. Edinburgh Sect. A, Vol. 125, January 1995, 13-29.

[10] J.R. Ringrose, *Exponential length and exponential rank in C*-algebras*, Proc. London Math. Soc. (3) 46(1983), 301-333.

Current Address:

Kun Wang: Department of Mathematics, University of Puerto Rico, Rio Piedras Campus, P.O.Box 70377, San Juan, Puerto Rico, USA 00931.

Email: lingling-1106@163.com